

A general tensor product theorem

Anton Deitmar

Abstract: We show that every admissible irreducible representation of a product of two locally compact groups is a tensor product of admissible irreducible representations of the factors.

Introduction

In the classical case the tensor product theorem is formulated for either totally disconnected groups or Lie groups and the proof by D. Flath [2], which is found in the books until this day, uses different methods for each case. The tensor product theorem is used in the theory of automorphic forms, to prove that every irreducible admissible representation of an adèle group $G(\mathbb{A})$ is an infinite product of local representations. In this note we give a unified proof which is simpler and applies to all locally compact groups. The proof essentially contains no new ideas. We only streamline and unify the existing proofs by using a more universal concept of Hecke algebras. This proof will also appear in the upcoming book by the author on Automorphic Forms.

1 Preliminaries

For a locally compact group G we denote by $\hat{G}$ the *unitary dual*, i.e., the set of isomorphism classes of irreducible unitary representations of G .

For a representation (π, V_π) of the compact group K and an irreducible representation (τ, V_τ) of K we define the τ -*isotype* as the sum of all subrepresentations of π , which are isomorphic to τ . If π is unitary, then V_π is the direct sum of its isotypes $V_\pi = \bigoplus_{\tau \in \hat{K}} V_\pi(\tau)$.

For a given representation $\tau \in \hat{K}$ let

$$e_\tau(k) = (\dim \tau) \overline{\text{tr}(\tau(k))}, \quad k \in K.$$

Lemma 1.1 *The function e_τ is an idempotent in the convolution algebra $C(K)$, i.e., one has $e_\tau * e_\tau = e_\tau$. Let (π, V_π) be a representation of K . Then the map*

$$P_\tau = \int_K e_\tau(k) \pi(k) dk$$

is a projection onto the isotype $V_\pi(\tau)$. Here we have normalized the Haar measure on K so that $\text{vol}(K) = 1$. If π is unitary, then P_τ is the orthogonal projection onto $V_\pi(\tau)$.

*If γ is another irreducible unitary representation of K which is not isomorphic to τ , then $e_\tau * e_\gamma = 0$.*

Proof: This is a direct consequence of the Peter-Weyl Theorem [1]. □

Let $F \subset \hat{K}$ be a finite subset. Then the function

$$e_F \stackrel{\text{def}}{=} \sum_{\tau \in F} e_\tau$$

is an idempotent in $C(K)$, because

$$e_F * e_F = \sum_{\tau, \gamma \in F} e_\tau * e_\gamma = \sum_{\tau \in F} e_\tau = e_F.$$

2 Main Theorem

Let G be a locally-compact group and $K \subset G$ a compact subgroup. A unitary representation (π, V_π) of G is called *K-admissible*, if the representation $\pi|_K$ decomposes with finite multiplicities, i.e., if for every $\tau \in \hat{K}$ the multiplicity

$$[\pi : \tau] = \dim \text{Hom}_K(V_\tau, V_\pi) = \dim \text{Hom}_K(V_\pi, V_\tau)$$

is finite. The representation π is *K-admissible* if and only if every *K*-isotype $V_\pi(\tau)$ is finite-dimensional. We denote by $\hat{G}_K$ the set of all isomorphism classes of irreducible unitary G -representations which are *K-admissible*. A representation π of G is called *admissible*, if there is a compact subgroup K such that π is *K-admissible*. We write $\hat{G}_{\text{adm}}$ for the set of isomorphism classes of irreducible unitary admissible representations of G .

Theorem 2.1 (Main Theorem) *Let G and H be locally compact groups. For any two $\pi \in \hat{G}_{\text{adm}}$ and $\eta \in \hat{H}_{\text{adm}}$ the tensor product $\pi \otimes \eta$ is in $\widehat{G \times H}_{\text{adm}}$ and the tensor map $(\pi, \eta) \mapsto \pi \otimes \eta$ is a bijection*

$$\hat{G}_{\text{adm}} \times \hat{H}_{\text{adm}} \xrightarrow{\cong} \widehat{G \times H}_{\text{adm}}.$$

More precisely, given two compact subgroups $K \subset G$ and $L \subset H$, the tensor map map is a bijection $\widehat{G}_K \times \widehat{H}_L \rightarrow \widehat{G \times H}_{K \times L}$.

The injectivity of the tensor map follows from the next lemma.

Lemma 2.2 *Let G be a locally compact group and let (η, V_η) be an irreducible unitary representation. Let W be a Hilbert space. Then every closed G -stable subspace of $V_\eta \otimes W$ is of the form $V_\eta \otimes W_1$ for a closed subspace W_1 of W . Here the tensor product refers to the tensor product in the category of Hilbert spaces, i.e., the Hilbert-space completion of the algebraic tensor product.*

In particular, it follows that for two locally compact groups G, H and irreducible unitary representations π, τ of G and H respectively, the representation $\pi \otimes \tau$ of $G \times H$ is irreducible and that $\pi \otimes \tau \cong \pi' \otimes \tau'$ implies $\pi \cong \pi'$ and $\tau \cong \tau'$.

Proof: Let $U \subset V_\eta \otimes W$ be a closed, G -stable subspace. Then the orthogonal projection P with image U is a bounded operator on $V_\eta \otimes W$ which commutes with all $\eta(g)$, $g \in G$. We show that every such operator T is of the form $1 \otimes S$ for an operator $S \in \mathcal{B}(W)$. So let T be a bounded operator on $V_\eta \otimes W$ with

$$T(\eta(g) \otimes 1) = (\eta(g) \otimes 1)T$$

for every $g \in G$. Let $(e_i)_{i \in I}$ be an orthonormal basis of W and for $j \in I$ let P_j be the orthogonal projection onto the subspace $\mathbb{C}e_j$. For $i, j \in I$ consider the operator $T_{i,j}$ given as composition:

$$V_\eta \rightarrow V_\eta \otimes e_i \xrightarrow{T} V_\eta \otimes W \xrightarrow{1 \otimes P_j} V_\eta \otimes e_j \rightarrow V_\eta.$$

This operator is bounded and commutes with the G -action. By the Lemma of Schur there exists a complex number $a_{i,j}$ with $T_{i,j} = a_{i,j}\text{Id}$. Since $\sum_j P_j = \text{Id}_W$, we get

$$T(v \otimes e_i) = \sum_j a_{i,j} v \otimes e_j = v \otimes S(e_i),$$

where $S(e_i) = \sum_j a_{i,j} e_j$ and the sum converges in W . As T is continuous, S extends to a unique continuous operator $S : W \rightarrow W$, satisfying

$$T(v \otimes w) = v \otimes S(w)$$

for all $v \in V_\eta$, $w \in W$. This means $T = 1 \otimes S$, as claimed.

We can apply this to the orthogonal projection P onto the invariant subspace U , which therefore is of the form $P = 1 \otimes P_1$. Then P_1 is again an orthogonal projection. Let W_1 be the image of P_1 , then $U = V_\eta \otimes W_1$.

We next show the irreducibility of $\pi \otimes \tau$. The first part implies that a closed, $G \times H$ -stable subspace U has to be of the form $U = V_\pi \otimes U_\tau$ as well as of the form $U = U_\pi \otimes V_\tau$ with closed subspaces $U_\pi \subset V_\pi$ and $U_\tau \subset V_\tau$. These subspaces have to be invariant themselves, hence the full spaces.

Finally let $\pi \otimes \tau \cong \pi' \otimes \tau'$, so there is a unitary isomorphism $T : V_\pi \otimes V_\tau \rightarrow V_{\pi'} \otimes V_{\tau'}$ commuting with the $G \times H$ -operation. Choose $w \in V_\tau$ with $\|w\| = 1$. Then $V_\pi \otimes w$ is an irreducible G -subrepresentation, therefore there is $w' \in V_{\tau'}$ mit $\|w'\| = 1$, such that $T(V_\pi \otimes w) = V_{\pi'} \otimes w'$. The map

$$V_\pi \rightarrow V_\pi \otimes w \xrightarrow{T} V_{\pi'} \otimes w' \rightarrow V_{\pi'}$$

is a unitary G -isomorphism, so we get $\pi \cong \pi'$ and analogously $\tau \cong \tau'$. $\square$

3 The Hecke algebra

Let G be a locally compact group. The set $C_c(G)$ of all continuous functions of compact support is a complex algebra under the convolution product

$$f * g(x) = \int_G f(y)g(y^{-1}x) dy = \int_G f(xy)g(y^{-1}) dy, \quad f, g \in C_c(G).$$

Let $K \subset G$ be a compact subgroup. For $\alpha, \beta \in C(K)$, there is the K -convolution

$$\alpha * \beta(k) = \int_K \alpha(l)\beta(l^{-1}k) dl$$

making $C(K)$ a $\mathbb{C}$ -algebra as well. We can also define a convolution between functions on G and K as follows. For $\alpha \in C(K)$ and $f \in C_c(G)$ we define

$$\alpha * f(x) = \int_K \alpha(k)f(k^{-1}x) dk \quad \text{and} \quad f * \alpha(x) = \int_K f(xk)\alpha(k^{-1}) dk.$$

We normalize the Haar-measure of the compact group K so that $\text{vol}(K) = 1$. If K is not a nullset in G , we insist that the Haar measure of G be normalized also to give K the volume 1. Hence in this case the Haar measures of G and K coincide on K .

Lemma 3.1 *With this normalization the convolution products are compatible in the sense that*

$$f * (g * h) = (f * g) * h.$$

*holds for all combinations of f, g, h being functions in either $C_c(G)$ or $C(K)$. Also we always have $(f * g)^* = g^* * f^*$, where $f^*(x) = \Delta(x^{-1})\overline{f(x^{-1})}$ and $\Delta(x)$ is the modular function of G .*

Proof: A computation. □

Let G be a locally compact group with a compact subgroup K . A function f on G is called K -finite, if the set of all functions $x \mapsto f(k_1 x k_2)$, $k_1, k_2 \in K$ spans a finite-dimensional vector space. It is a simple observation that the convolution product of two K -finite functions is K -finite. We define the *Hecke-algebra* $\mathcal{H} = \mathcal{H}_{G,K}$ of the pair (G, K) as the convolution algebra of all K -finite functions in $C_c(G)$.

Lemma 3.2 *Let G be a locally compact group and K a compact subgroup.*

(a) *Let I_K be the set of all finite subsets of $\hat{K}$. For $F \in I_K$ set $e_F = \sum_{\tau \in F} e_\tau$ and*

$$C_F \stackrel{\text{def}}{=} e_F * C_c(G) * e_F = e_F * \mathcal{H} * e_F.$$

Then e_F is an idempotent in $C(K)$ and C_F is a subalgebra of $\mathcal{H}$. The Hecke-algebra $\mathcal{H}$ is the union of all these subalgebras.

If $F \subset F'$ in I_K , then $C_F \subset C_{F'}$. If $F = \{\tau\}$ we also write $C_F = C_\tau$.

(b) *For a unitary representation (π, V_π) of G the space $\pi(\mathcal{H})V_\pi$ is dense in V_π .*

(c) *The Hecke-algebra is a $*$ -Algebra with involution $f^*(x) = \Delta(x^{-1})\overline{f(x^{-1})}$. If π is a unitary representation of G , then π defines a $*$ -representation of $\mathcal{H}$. For two unitary representations π, π' of G we have*

$$\pi \cong_G \pi' \Leftrightarrow \pi \cong_{\mathcal{H}} \pi',$$

so, π and π' are unitarily equivalent if and only if they define isomorphic $\mathcal{H}$ -modules.

Proof: Part (a) is clear. We show (b). For $h \in C(K)$ we write

$$\pi(h) = \int_K h(k) \pi(k) dk.$$

Let $F \in I_K$ and $P_F = \pi(e_F)$. Then $P_F^2 = \pi(e_F)\pi(e_F) = \pi(e_F * e_F) = \pi(e_F) = P_F$. So P_F is a projection with image $V_\pi(F) = \bigoplus_{\tau \in F} V_\pi(\tau)$, which is the F -isotype of π and the kernel is $\bigoplus_{\tau \notin F} V_\pi(\tau)$. Hence the union of all $V_\pi(F)$ with $F \in I_K$ is dense in V_π . So it suffices to show that $\pi(C_F)V_\pi$ is dense in $V_\pi(F)$. Let $v \in V_\pi(F)$ and let $\varepsilon > 0$. Since $\pi(e_F)$ is continuous, there is $C > 0$ such that $\|\pi(e_F)w\| \leq C\|w\|$ for every $w \in V_\pi$. Since the map $G \times V_\pi \rightarrow V_\pi$; $(g, v) \mapsto \pi(g)v$ is continuous, there is a neighborhood U of the unit in G , such that $x \in U \Rightarrow \|\pi(x)v - v\| < \varepsilon/C$. Let $f \in C_c(G)$ with support in U such that $f \geq 0$ and $\int_G f(x) dx = 1$. Then

$$\|\pi(f)v - v\| = \left\| \int_G f(x) (\pi(x)v - v) dx \right\| \leq \int_G f(x) \|\pi(x)v - v\| dx < \varepsilon/C.$$

One has $e_F * f * e_F \in C_F \subset \mathcal{H}$ and

$$\|\pi(e_F * f * e_F)v - v\| = \|\pi(e_F)(\pi(f)v - v)\| < \varepsilon.$$

This shows (b). We continue with (c). The closedness under $*$ is clear. Let π be a unitary G -representation, then for $f \in \mathcal{H}$ one has

$$\begin{aligned} \pi(f^*) &= \int_G \Delta(x^{-1}) \overline{f(x^{-1})} \pi(x) dx = \int_G \overline{f(x)} \pi(x^{-1}) dx \\ &= \int_G \overline{f(x)} \pi(x)^* dx = \left(\int_G f(x) \pi(x) dx \right)^* = \pi(f)^*. \end{aligned}$$

If π and π' are isomorphic as G -representations, then so they are as $\mathcal{H}$ -modules. For the converse let $T : V_\pi \rightarrow V_{\pi'}$ be a unitary $\mathcal{H}$ -isomorphism, so $T\pi(f) = \pi'(f)T$ for every $f \in \mathcal{H}$. We first remark, that the same must hold for $f \in C_c(G)$. For this let $S = T\pi(f) - \pi'(f)T$, then for every finite subset F of $\hat{K}$,

$$\pi'(e_F)S\pi(e_F) = T\pi(\underbrace{e_F f e_F}_{\in \mathcal{H}}) - \pi'(e_F f e_F)T = 0.$$

Therefore $Sv = 0$ for every vector $v \in V_\pi(F)$ and since the $V_\pi(F)$ span a dense subspace of V_π , we get $S = 0$, so $T\pi(f) = \pi'(f)T$ for every $f \in C_c(G)$.

Let $\varepsilon > 0$ and let $x \in G$, $v \in V$. Then, by the continuity of the representations π and π' there exists a neighborhood U of $x \in G$ such that $\|\pi(u)v - \pi(x)v\| < \varepsilon/2$ and $\|\pi'(u)Tv - \pi'(x)Tv\| < \varepsilon/2$. Let $f \in C_c(G)$ have support inside U and suppose $f \geq 0$ and $\int_G f(x) dx = 1$. Since T is unitary, we have

$$\begin{aligned} \|T\pi(f)v - T\pi(x)v\| &= \|\pi(f)v - \pi(x)v\| \\ &= \left\| \int_G f(u)(\pi(u)v - \pi(x)v) du \right\| \leq \int_G f(u)\|\pi(u)v - \pi(x)v\| du < \varepsilon/2 \end{aligned}$$

and likewise $\|T\pi(f)v - \pi'(x)Tv\| = \|\pi'(f)Tv - \pi'(x)Tv\| < \varepsilon/2$. This implies $\|T\pi(x)v - \pi'(x)Tv\| < \varepsilon$. Since $\varepsilon > 0$ and $v \in V_\pi$ are arbitrary, we conclude $T\pi(x) = \pi'(x)T$. $\square$

If (π, V_π) is an irreducible unitary representation of G , then every $f \in C_c(G)$ defines a continuous linear operator $\pi(f)$ on V_π . For $0 \neq v \in V_\pi$ the space $\pi(C_c(G))v$ is a G -stable subspace of V_π . As π is irreducible, the space $\pi(\mathcal{H})v$ is a dense subspace.

Proposition 3.3 *Let G be a locally compact group and $K \subset G$ a compact subgroup.*

- (a) *Let (π, V_π) be an irreducible representation of G . Let F be a finite subset of $\hat{K}$. Then the F -isotype F -Isotyp $V_\pi(F) = \bigoplus_{\tau \in F} V_\pi(\tau)$ is an irreducible C_F -module. In particular, $\pi(\mathcal{H})V_\pi$ is an irreducible $\mathcal{H}$ -module.*
- (b) *Let (η, V_η) be a unitary representation of G and F a finite subset of $\hat{K}$. If M is a finite-dimensional irreducible C_F -submodule of $V_\eta(F)$, then the G -subrepresentation of π generated by M is irreducible.*
- (c) *Let η, π be irreducible unitary representations of G , which are K -admissible. Then*

$$\eta \cong \pi \Leftrightarrow V_\eta(F) \cong V_\pi(F) \text{ for every } F \in I_K,$$

where on the right we consider isomorphism as C_F -modules.

Proof: (a) To show irreducibility of $V_\pi(F)$, let $0 \neq U \subset V_\pi(F)$ be a closed submodule. Since V_π is irreducible, the space $\pi(C_c(G))U$ is dense in V_π . Let $P_F : V_\pi \rightarrow V_\pi(F)$ be the isotypical projection. For $h \in C(K)$ we write

$$\pi(h) = \int_K h(k)\pi(k) dk.$$

Then we have $P_F = \pi(e_F)$. Let $f \in C_c(G)$. Then

$$\pi(f)\pi(e_\tau) = \pi(f * e_\tau) \quad \text{und} \quad \pi(e_\tau)\pi(f) = \pi(e_\tau * f),$$

as a calculation shows.

The projection P_F is continuous, so the space $P_F(C_c(G)U)$ is dense in $V_\pi(F)$, therefore $\overline{P_F(C_c(G)U)} = V_\pi(F)$. Since $P_F = \pi(e_F)$ we get

$$\begin{aligned} V_\pi(F) &= \overline{P_F(C_c(G)U)} = \overline{\pi(e_F)C_c(G)U} \\ &= \overline{\pi(e_F * C_c(G) * e_F)U} = \overline{\pi(C_F)U} = \overline{U} = U, \end{aligned}$$

which shows that $V_\pi(F)$ is irreducible.

We now show (b). Let $m_0 \in M \setminus \{0\}$ and let $\text{Ann}(m_0) = \{a \in C_F : am_0 = 0\}$ be its annihilator. Then $\text{Ann}(m_0)$ is a left ideal of C_F and the map $a \mapsto am_0$ is a module isomorphism

$$C_F / \text{Ann}(m_0) \rightarrow C_F m_0.$$

Since M is finite-dimensional and irreducible it follows $M = C_F m_0$, so $M \cong C_F / J$ with $J = \text{Ann}(m_0)$. Let U be the G -stable closed subspace of V_π generated by M . We show $P_F(U) = M$, where $P_F = \pi(e_F)$ is the orthogonal projection onto M .

Claim: Let $\overline{J}$ be the annihilator $\text{Ann}_{C_c(G)}(m_0)$ of m_0 in $C_c(G)$. Then

$$\overline{J} = \overline{J}e_F \oplus \text{Ann}_{C_c(G)}(e_F),$$

where $\text{Ann}_{C_c(G)}(e_F)$ is the annihilator of e_F in $C_c(G)$, which is the set of all $f \in C_c(G)$ with $f * e_F = 0$.

We show the claim. It is clear that $\overline{J}e_F \subset \text{Ann}_{C_c(G)}(v_0) = \overline{J}$. Further $v_0 = e_F v_0$ and so $\text{Ann}_{C_c(G)}(e_F) \subset \text{Ann}_{C_c(G)}(v_0) = \overline{J}$. It remains to show that $\overline{J}$ lies in the right hand side. Since e_F is an idempotent, we have $C_c(G) = C_c(G)e_F \oplus \text{Ann}_{C_c(G)}(e_F)$, since every $f \in C_c(G)$ can be written as $f = fe_F + (f - fe_F)$ and $f - fe_F$ lies in the annihilator of e_F . Further $\text{Ann}_{C_c(G)}(e_F) \subset \text{Ann}_{C_c(G)}(v_0) = \overline{J}$, which shows the claim.

So we have

$$C_c(G)v_0 \cong C_c(G)/\overline{J} \cong C_c(G)e_F/(\overline{J}e_F).$$

This implies

$$P_F(C_c(G)v_0) \cong e_F C_c(G)e_F/(\overline{J}e_F) \cong C_F/J \cong M.$$

Therefore $P_F(U) = M$. Let now U' be a subrepresentation of U . Then $P_F(U') = 0$ or M . In the first case $M \subset (U')^\perp$ and so $U \subset (U')^\perp$, since the space $(U')^\perp$ is a subrepresentation. This gives $U' = 0$. In the second case $M \subset U'$ and so $U' = U$. Indeed we conclude that U is irreducible.

(c) Let $(\eta, V_\eta), (\pi, V_\pi)$ be isomorphic representations, then so are the C_F -modules $V_\eta(F)$ and $V_\pi(F)$. For the converse let for every F be given a C_F -isomorphism $\phi_F : V_\eta(F) \rightarrow V_\pi(F)$. According to the Lemma of Schur two isomorphisms for a given F only differ by a multiple scalar. So we can normalize all these isomorphisms in a manner that they extend each other, i.e., that we have $\phi_F = \phi_{F'}|_{V_\eta(F)}$, if $F \subset F'$. This means that the ϕ_F can be put together to a $\mathcal{H}$ -isomorphism

$$\phi : \eta(\mathcal{H})V_\eta \xrightarrow{\cong} \pi(\mathcal{H})V_\pi.$$

Both sides are dense sub vector spaces. If we can show that ϕ is isometric, this map extends to an isomorphism of G -representations by Lemma 3.2. Let $0 \neq v_0 \in \eta(\mathcal{H})V_\eta$ and let $w_0 = \phi(v_0)$. We can assume that $\|v_0\| = \|w_0\| = 1$ and we claim that ϕ is isometric. For this we transport both inner products to the same side via ϕ , so we have two inner products $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$ on, say $\pi(\mathcal{H})V_\pi$ such that v_0 has norm 1 in both inner products and the operation of $\mathcal{H}$ is a $*$ -operation in both inner products. On the finite dimensional space $\pi(C_F)V_\pi$ these inner products must coincide by the Lemma of Schur. This holds for every $F \in I_K$, so that ϕ is indeed isometric. $\square$

Let now G, H locally compact with compact subgroups $K \subset G$ and $L \subset H$. Let E be a finite subset of $\hat{K}$ and F a finite subset of $\hat{L}$. There is a natural Homomorphism

$$\psi : C_c(G) \otimes C_c(H) \rightarrow G_c(G \times H)$$

given by

$$\psi(f \otimes g)(x, y) = f(x)g(y).$$

This induces a homomorphism

$$C_E \otimes C_F \rightarrow C_{E \times F}.$$

This map is not surjective in general. But this fact causes no harm, as we have

Lemma 3.4 *If M is a finite-dimensional irreducible $C_{E \times F}$ -*-submodule of a unitary G -representation, then it is irreducible as $C_E \otimes C_F$ -module.*

Proof: We equip $C_{E \times F} \subset L^1(G \times H)$ with the topology of the L^1 -norm. Then $C_E \otimes C_F$ is dense in $C_{E \times F}$, as $C_c(G)$ is dense in $L^1(G)$. The representation $\rho : C_{E \times F} \rightarrow \text{End}_{\mathbb{C}}(M)$ is continuous, since M comes from a unitary representation of $G \times H$. So the image of $C_E \otimes C_F$ is dense in the (finite-dimensional) image of $C_{E \times F}$ in $\text{End}(M)$, so these two images agree. $\square$

In order to show Theorem 2.1 we only need the second assertion of the following lemma.

Lemma 3.5 (a) *Let A be a $\mathbb{C}$ -algebra with unit and M a simple A -module, which is finite dimensional over $\mathbb{C}$. Then the map $A \rightarrow \text{End}_{\mathbb{C}}(M)$ is surjective.*

(b) *Let A, B complex algebras with units and let $R = A \otimes B$. If M, N are simple modules over A and B resp., which are finite-dimensional over $\mathbb{C}$, then $M \otimes N$ is a simple R -module and every simple R -module, which is finite-dimensional over $\mathbb{C}$, is of this form with uniquely determined modules M and N .*

Proof: Part (a) is a well-known result of Wedderburn. See [3] for a proof.

We show (b). According to (a) it suffices to assume $A = \text{End}_{\mathbb{C}}(M)$ and $B = \text{End}_{\mathbb{C}}(N)$ for the first assertion. The canonical map from $\text{End}_{\mathbb{C}}(M) \otimes \text{End}_{\mathbb{C}}(N)$ to $\text{End}_{\mathbb{C}}(M \otimes N)$ is surjective, and so $M \otimes N$ is simple.

Let now V be a given $\mathbb{C}$ -finite-dimensional simple $A \otimes B$ -module. Then V contains a simple A -module M as V is finite-dimensional. Let $N = \text{Hom}_A(M, V)$. This vector space is a B -module in the obvious way. Consider the map $\phi : M \otimes N \rightarrow V$ given by

$$\phi(m \otimes \alpha) = \alpha(m).$$

Then $\phi \neq 0$ is an $A \otimes B$ -homomorphism, hence surjective, as V is simple. We have to show that ϕ has zero kernel. For this let $\alpha_1, \dots, \alpha_k$ be a basis of N and $m_1, \dots, m_l$ a basis of M . Let $c_{i,j} \in \mathbb{C}$ be given with $\phi\left(\sum_{i,j} c_{i,j} m_i \otimes \alpha_j\right) = 0$. We have to show that all coefficients $c_{i,j}$ are zero. We have

$$0 = \phi\left(\sum_{i,j} c_{i,j} m_i \otimes \alpha_j\right) = \sum_{i,j} c_{i,j} \alpha_j(m_i) = \sum_j \alpha_j\left(\sum_i c_{i,j} m_i\right).$$

Let $P : M \rightarrow M$ be a projection onto a one-dimensional subspace, say $\mathbb{C}m_0$. By part (a) there is $a \in A$ with $am = Pm$ for every $m \in M$. We conclude

$$0 = \sum_j \alpha_j\left(P\left(\sum_i c_{i,j} m_i\right)\right) = \sum_j \lambda_j \alpha_j(m_0),$$

where $P(\sum_i c_{i,j} m_i) = \lambda_j m_0$. For $a \in A$ arbitrary, we get $0 = \sum_j \lambda_j \alpha_j(am_0)$. As a runs through A , the vector am_0 runs through M , so $\sum_j \lambda_j \alpha_j = 0$. Since the α_j are linearly independent, all λ_j are zero, hence all $\sum_i c_{i,j} m_i$ are zero and again by linear independence, we get $c_{i,j} = 0$. This proof also shows that all simple A -submodules of V are isomorphic, whence the uniqueness. $\square$

We now prove Theorem 2.1. Let η be a $K \times L$ -admissible representation of $G \times H$. Let E be a finite subset of $\hat{K}$ and F a finite subset of $\hat{L}$. Then $V_{\eta}(E \times F)$ is a finite-dimensional irreducible $C_{E \times F}$ -module. By Lemma 3.4 the space $V_{\eta}(E \times F)$ is an irreducible $C_E \otimes C_F$ -module and by Lemma 3.5 the module $V_{\eta}(E \times F)$ is a tensor product of modules, which we write as $V_{\pi}(E) \otimes V_{\tau}(F)$. The uniqueness of the

tensor-factors give injective homomorphisms $\varphi_E^{E'} : V_\pi(E) \rightarrow V_\pi(E')$ if $E \subset E'$ and likewise for F . By the uniqueness of the inner products in the Lemma of Schur, we can scale these homomorphisms in a way that they are isometric. We define

$$\tilde{V}_\pi \stackrel{\text{def}}{=} \lim_{\rightarrow} V_\pi(E).$$

Then $\tilde{V}_\pi$ is a pre-Hilbert-space and we write V_π for its completion. Analogously we construct V_τ . For every finite subset $E \subset \hat{K}$ the algebra C_E acts on $\tilde{V}_\pi$ by continuous operators. These extend continuously to V_π and we get $*$ -representation of the Hecke-algebra $\mathcal{H}_G$ and likewise for V_τ . By construction the isotypes of V_π are precisely the spaces $V_\pi(E)$ for $E \in I_K$. The isometrical maps $V_\pi(E) \otimes V_\tau(F) \hookrightarrow V_\eta$ induce an isometric $\mathcal{H}_G \otimes \mathcal{H}_H$ -homomorphism $\Phi : V_\pi \otimes V_\tau \rightarrow V_\eta$, which by irreducibility must be an isomorphism. It only remains to install a unitary representation of the group $G \times H$ on $V_\pi \otimes V_\tau$, so at first a representation π of G on V_π . Fix an arbitrary vector $w \in V_\tau$ with $\|w\| = 1$. Then the map $V_\pi \rightarrow V_\pi \otimes w \xrightarrow{\Phi} V_\eta$ is an isometric embedding of V_π into V_η , which commutes with the $\mathcal{H}_G$ -operation. The G -representation on V_η then defines a unitary G -representation on V_π , which induces the $\mathcal{H}_G$ -representation. We do the same with H and get irreducible unitary representations π and τ of G and H such that Φ is a $G \times H$ -isomorphism. $\square$

References

- [1] **Deitmar, Anton; Echterhoff, Siegfried:** *Principles of harmonic analysis*. Universitext. Springer, New York, 2009.
- [2] **Flath, D.:** *Decomposition of representations into tensor products*. Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1, pp. 179–183, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979.
- [3] **Lang, Serge:** *Algebra*. Revised third edition. Graduate Texts in Mathematics, 211. Springer-Verlag, New York, 2002

Mathematisches Institut, Auf der Morgenstelle 10, 72076 Tübingen, Germany
 deitmar@uni-tuebingen.de